

C3, IYGB, PAPER V

1.

$$f(x) = x \ln(1+x^2)$$

• BY PRODUCT RULE

$$f'(x) = 1 \times \ln(1+x^2) + x \times \frac{1}{1+x^2} \times 2x$$

$$f'(x) = \ln(1+x^2) + \frac{2x^2}{1+x^2}$$

$$f'(1) = \ln 2 + \frac{2}{2}$$

$$\therefore \text{GRADIENT } 1 + \ln 2$$

• $f(1) = \ln 2 \quad \therefore (1, \ln 2)$

TANGENT EQUATION

$$y - y_0 = m(x - x_0)$$

$$y - \ln 2 = (1 + \ln 2)(x - 1)$$

$$y - \ln 2 = (1 + \ln 2)x - (1 + \ln 2)$$


$$y - \ln 2 = x(1 + \ln 2) - 1 - \ln 2$$

$$y = x(1 + \ln 2) - 1$$




AS REQUIRED

2.

$$\sin x = \frac{12}{13}$$



$$\cos y = \frac{15}{17}$$



$$\sin(x-y) = \sin x \cos y - \cos x \sin y$$

$$= \frac{12}{13} \times \frac{15}{17} - \left(-\frac{5}{13}\right) \frac{8}{17}$$

$$= \frac{180}{221} + \frac{40}{221}$$

$$= \frac{220}{221}$$

C3, NYGB, PAPER V

-2-

3.

$$\begin{cases} f(x) = x^2 + 3x - 7, & x \in \mathbb{R} \\ g(x) = ax + b, & x \in \mathbb{R} \end{cases}$$

$$f(g(x)) = f(ax+b) = (ax+b)^2 + 3(ax+b) - 7$$

$$g(f(x)) = g(x^2 + 3x - 7) = a(x^2 + 3x - 7) + b$$

$$\textcircled{\ast} \quad f(g(-2)) = 21$$

$$\textcircled{\ast} \quad g(f(1)) = 0$$

$$(-2a+b)^2 + 3(-2a+b) - 7 = 21$$

$$a(1+3-7) + b = 0$$

$$-3a + b = 0$$

$$\boxed{b = 3a}$$

$$(-2a+3a)^2 + 3(-2a+3a) - 28 = 0$$

$$a^2 + 3a - 28 = 0$$

$$(a+7)(a-4) = 0$$

$$a = \begin{cases} 4 \\ -7 \end{cases}$$

$$b = 12$$



4. a)

$$\begin{cases} f(x) = a \ln bx, & x > 0 \end{cases}$$

$$\textcircled{\ast} \quad \left(\frac{1}{3}\right)^0 \Rightarrow 0 = a \ln\left(\frac{1}{3}b\right)$$

$$\Rightarrow \ln\left(\frac{1}{3}b\right) = 0 \quad (a \neq 0)$$

$$\Rightarrow \frac{1}{3}b = e^0$$

$$\Rightarrow \frac{1}{3}b = 1$$

$$\Rightarrow b = 3$$




$$\textcircled{a} (3, 4) \Rightarrow 4 = a \ln(3x3)$$

$$\Rightarrow 4 = a \ln 9$$

$$\Rightarrow 4 = 2a \ln 3$$

$$\Rightarrow a = \frac{2}{\ln 3}$$

b)



c) $\left\{ \begin{array}{l} |f(x)| = 8 \\ \end{array} \right.$ DEFINITELY HAS TWO SOLUTIONS (from PAPER V)

$$\textcircled{1} \frac{2}{\ln 3} \ln(3x) = 8$$

$$\Rightarrow \ln 3x = 4 \ln 3$$

$$\Rightarrow \ln 3x = \ln 81$$

$$\Rightarrow 3x = 81$$

$$\Rightarrow x = 27$$

$$\textcircled{2} \frac{2}{\ln 3} \ln 3x = -8$$

$$\ln 3x = -4 \ln 3$$

$$\ln 3x = \ln \left(\frac{1}{81}\right)$$

$$3x = \frac{1}{81}$$

$$x = \frac{1}{243}$$

$$\begin{aligned}
 5. \text{ a) } \cos 3x &= \cos(2x+x) \\
 &= \cos 2x \cos x - \sin 2x \sin x \\
 &= (2\cos^2 x - 1) \cos x - (2\sin x \cos x) \sin x \\
 &= 2\cos^3 x - \cos x - 2\sin^2 x \cos x \\
 &= 2\cos^3 x - \cos x - 2(1 - \cos^2 x) \cos x \\
 &= 2\cos^3 x - \cos x - 2\cos x + 2\cos^3 x \\
 &= 4\cos^3 x - 3\cos x
 \end{aligned}$$

~~AS FÜR IRAD~~

b) $8\cos^3 x - 6\cos x + 1 = 0$

$$\begin{aligned}
 8\cos^3 x - 6\cos x &= -1 \\
 4\cos^3 x - 3\cos x &= -\frac{1}{2} \quad) \div 2
 \end{aligned}$$

$$\cos 3x = -\frac{1}{2}$$

$$\arccos\left(-\frac{1}{2}\right) = \frac{2\pi}{3}$$

$$\begin{cases} 3x = \frac{2\pi}{3} \pm 2n\pi \\ 3x = \frac{4\pi}{3} \pm 2n\pi \end{cases} \quad n=0,1,2,3,\dots$$

$$\begin{cases} x = \frac{2\pi}{9} \pm \frac{2n\pi}{3} \\ x = \frac{4\pi}{9} \pm \frac{2n\pi}{3} \end{cases}$$

$$x = \frac{2\pi}{9}, \frac{8\pi}{9}, \frac{14\pi}{9}, \frac{4\pi}{9}, \frac{10\pi}{9}, \frac{16\pi}{9}$$

~~AS FÜR IRAD~~

6. a)

$$y = 4x^{\frac{3}{2}} - \frac{7}{8} \ln 4x$$

$$\frac{dy}{dx} = 6x^{\frac{1}{2}} - \frac{7}{8} \times \frac{1}{x}$$

$$\left. \frac{dy}{dx} \right|_{x=1/4} = 6 \times \frac{1}{2} - \frac{7}{8} \times 4 = 3 - \frac{7}{2} = -\frac{1}{2}$$

• Normal gradient = 2

• $y = 4 \times \left(\frac{1}{4}\right)^{\frac{3}{2}} - \cancel{\frac{7}{8} \ln 1} = \frac{1}{2}$ at $(\frac{1}{4}, \frac{1}{2})$

$$y - y_0 = m(x - x_0)$$

$$y - \frac{1}{2} = 2(x - \frac{1}{4})$$

$$y - \frac{1}{2} = 2x - \frac{1}{2}$$

$$\cancel{y = 2x}$$

b)

$$y = 2x \quad \& \quad y = 4x^{\frac{3}{2}} - \frac{7}{8} \ln 4x$$

$$2x = 4x^{\frac{3}{2}} - \frac{7}{8} \ln 4x$$

$$2x + \frac{7}{8} \ln 4x = 4x^{\frac{3}{2}}$$

$$\frac{16x + 7 \ln 4x}{8} = 4x^{\frac{3}{2}}$$

$$x^{\frac{3}{2}} = \frac{16x + 7 \ln 4x}{32}$$

$$x = \left(\frac{16x + 7 \ln 4x}{32} \right)^{\frac{2}{3}} \cancel{\checkmark}$$

c)

$$x_{n+1} = \left[\frac{16x_n + 7 \ln(4x_n)}{32} \right]^{\frac{2}{3}}$$

$$x_0 = 0.7$$

$$x_1 \approx 0.692$$


$$x_2 \approx 0.686$$

$$x_3 \approx 0.683$$

$$x_4 \approx 0.680$$



d)



EQUATION SOLVED

$$2x = 4x^{\frac{3}{2}} - \frac{7}{8} \ln 4x$$

$$2x - 4x^{\frac{3}{2}} + \frac{7}{8} \ln 4x = 0$$

$$\text{Let } f(x) = 2x - 4x^{\frac{3}{2}} + \frac{7}{8} \ln 4x$$

$$\begin{aligned} f(0.67545) &= 0.000083 > 0 \\ f(0.67555) &= -0.00008 < 0 \end{aligned} \quad \Rightarrow$$

CHANGE OF SIGN (AND
CONTINUITY) IMPLY

$$0.67545 < \alpha < 0.6755$$

$$\alpha \approx 0.6755$$

CORRECT TO 4 d.p.

7.

$$y = \left(\frac{1-x}{1+x} \right)^{\frac{1}{2}} (2x+1) \quad -1 < x \leq 1$$

$$\Rightarrow \ln y = \ln \left[\left(\frac{1-x}{1+x} \right)^{\frac{1}{2}} (2x+1) \right]$$

$$\Rightarrow \ln y = \ln (1-x)^{\frac{1}{2}} - \ln (1+x)^{\frac{1}{2}} + \ln (2x+1)$$

$$\Rightarrow \ln y = \frac{1}{2} \ln (1-x) - \frac{1}{2} \ln (1+x) + \ln (2x+1)$$

~~Diff~~ w.r.t x

$$\Rightarrow \frac{1}{y} \frac{dy}{dx} = -\frac{1}{2} \times \frac{1}{1-x} - \frac{1}{2} \times \frac{1}{1+x} + \frac{1}{2x+1} \times 2$$

$$\Rightarrow \frac{dy}{dx} \frac{1}{y} = -\frac{2}{2x+1} - \frac{1}{2} \left(\frac{1}{1-x} + \frac{1}{1+x} \right)$$

$$\Rightarrow \frac{dy}{dx} \frac{1}{y} = -\frac{2}{2x+1} - \frac{1}{2} \left(\frac{1+x+1-x}{(1-x)(1+x)} \right)$$

$$\Rightarrow \frac{dy}{dx} \frac{1}{y} = -\frac{2}{2x+1} - \frac{1}{1-x^2}$$

$$\Rightarrow \frac{dy}{dx} = y \left[-\frac{2}{2x+1} - \frac{1}{1-x^2} \right]$$

$$\Rightarrow \frac{dy}{dx} = \left(\frac{1-x}{1+x} \right)^{\frac{1}{2}} (2x+1) \left[-\frac{2}{2x+1} - \frac{1}{1-x^2} \right]$$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{x=\frac{1}{2}} = \left(\frac{1}{2} \right)^{\frac{1}{2}} \times 2 \left[1 - \frac{1}{3/4} \right]$$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{x=\frac{1}{2}} = \left(\frac{1}{3} \right)^{\frac{1}{2}} \times 2 \left(-\frac{1}{3} \right) = -\frac{2}{3} \frac{1}{\sqrt{3}} = -\frac{2}{9} \sqrt{3}$$

C3, IYGB, PAPER V

8. a)

$$P_1 - P_2 = 4800 \text{ when } t=T$$

$$1600e^{\frac{1}{4}T} - 100e^{\frac{1}{2}T} = 4800$$

$$16e^{\frac{1}{4}T} - e^{\frac{1}{2}T} = 48$$

$$e^{\frac{1}{2}T} - 16e^{\frac{1}{4}T} + 48 = 0$$

$$(e^{\frac{1}{4}T} - 4)(e^{\frac{1}{4}T} - 12) = 0$$

$$e^{\frac{1}{4}T} = \begin{cases} 4 \\ 12 \end{cases}$$

$$\frac{1}{4}T = \begin{cases} \ln 4 \\ \ln 12 \end{cases}$$

$$T = \begin{cases} 4\ln 4 = 8\ln 2 \\ 4\ln 12 \end{cases}$$

b)

$$\text{LET } P = P_1 - P_2$$

$$P = 1600e^{\frac{1}{4}t} - 100e^{\frac{1}{2}t}$$

$$\frac{dP}{dt} = 400e^{\frac{1}{4}t} - 50e^{\frac{1}{2}t}$$

• solve for zero

$$\Rightarrow 400e^{\frac{1}{4}t} - 50e^{\frac{1}{2}t} = 0$$

$$\Rightarrow 400e^{\frac{1}{4}t} = 50e^{\frac{1}{2}t} \quad) \quad \div e^{\frac{1}{4}t}$$

$$\Rightarrow 8 = e^{\frac{1}{4}t} \quad) \quad \div 50$$

$$16 \boxed{e^{\frac{1}{4}t} = 8}$$

C3, IYGB, PAPER V

-9-

$$\therefore P = 1600e^{\frac{1}{4}t} - 100e^{\frac{1}{2}t}$$

$$\Rightarrow P = 1600(e^{\frac{1}{4}t}) - 100(e^{\frac{1}{4}t})^2$$

$$\Rightarrow P = 1600 \times 8 - 100 \times 8^2$$

$$\Rightarrow P = 12800 - 6400$$

$$\Rightarrow P = 6400$$

JUSTIFICATION NOT ACTUALLY NEEDED SINCE AT $t=T$, $P_1-P_2=4800$

$$\frac{d^2P}{dt^2} = 1600e^{\frac{1}{4}t} - 50e^{\frac{1}{2}t}$$

$$\left. \frac{d^2P}{dt^2} \right|_{e^{\frac{1}{4}t}=8} = 1600 \times 8 - 50 \times 8^2 = 800 - 3200 < 0$$

\therefore INDEFINITE