

IYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 1

MANIPULATE THE FUNCTION AS FOLLOWS

$$f(z) = \frac{1}{z+4} = \frac{1}{z(1+\frac{4}{z})} = \frac{1}{z} \left(1 + \frac{4}{z}\right)^{-1}$$

EXPANDING BINOMIALLY NOTING THAT THE RADIUS OF CONVERGENCE MUST BE

$$\left| \frac{4}{z} \right| < 1$$

$$\underline{|z| > 4}$$

THENCE WE OBTAIN

$$f(z) = \frac{1}{z} \left[1 - \frac{4}{z} + \frac{16}{z^2} - \frac{64}{z^3} + \frac{256}{z^4} - \dots \right]$$

$$f(z) = \frac{1}{z} - \frac{4}{z^2} + \frac{16}{z^3} - \frac{64}{z^4} + \frac{256}{z^5} - \dots$$


-1-

YGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 2

START BY USING THE CONTOUR $|z|=1$ OR $z = e^{i\theta}$ $0 \leq \theta < 2\pi$

$$\bullet \cos\theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta}) = \frac{1}{2}\left(z + \frac{1}{z}\right)$$

$$\begin{aligned}\bullet 4\cos\theta - 5 &= 4 \times \frac{1}{2}\left(z + \frac{1}{z}\right) - 5 \\ &= 2z + \frac{2}{z} - 5 \\ &= \frac{1}{z}(2z^2 - 5z + 2) \\ &= \frac{1}{z}(2z-1)(z-2)\end{aligned}$$



TRANSFORMING THE INTEGRAL

$$\begin{aligned}\int_0^{2\pi} \frac{1}{4\cos\theta - 5} d\theta &= \int_{\Gamma} \frac{1}{\frac{1}{z}(2z-1)(z-2)} \left(-\frac{i}{z} dz\right) \\ &= \int_{\Gamma} \frac{-i}{(2z-1)(z-2)} dz\end{aligned}$$

$$\begin{aligned}z &= e^{i\theta} \\ dz &= ie^{i\theta} d\theta \\ dz &= iz d\theta \\ d\theta &= \frac{dz}{iz} \\ d\theta &= -\frac{i}{z} dz\end{aligned}$$

NOW THE INTEGRAND HAS SIMPLE POLES AT

$z=2$ & $z=\frac{1}{2}$ OF WHICH ONLY THE ONE

AT $z=\frac{1}{2}$ IS INSIDE Γ - FIND RESIDUE

$$\begin{aligned}\lim_{z \rightarrow \frac{1}{2}} \left[\left(z - \frac{1}{2}\right) f(z) \right] &= \lim_{z \rightarrow \frac{1}{2}} \left[\left(z - \frac{1}{2}\right) \times \frac{-i}{(2z-1)(z-2)} \right] \\ &= \lim_{z \rightarrow \frac{1}{2}} \left[\cancel{\left(z - \frac{1}{2}\right)} \times \frac{-i}{2\cancel{\left(z - \frac{1}{2}\right)}(z-2)} \right] \\ &= \lim_{z \rightarrow \frac{1}{2}} \left[\frac{-i}{2(z-2)} \right] \\ &= \underline{\underline{\frac{1}{3}i}}\end{aligned}$$

IYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 2BY THE RESIDUE THEOREM WE HAVE

$$\int_{\Gamma} f(z) dz = 2\pi i \times \sum (\text{RESIDUES INSIDE } \Gamma)$$

$$\int_{\Gamma} \frac{-i}{(z-1)(z-2)} dz = 2\pi i \times \frac{1}{3}i$$

$$\int_0^{2\pi} \frac{1}{4\cos\theta - 5} d\theta = -\frac{2\pi}{3}$$

1YGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 3

Assume a solution of the form

$$y = \sum_{r=0}^{\infty} a_r x^{r+k}, \quad a_0 \neq 0, \quad k \in \mathbb{R}$$

$$\frac{dy}{dx} = \sum_{r=0}^{\infty} a_r (r+k) x^{r+k-1}$$

$$\frac{d^2y}{dx^2} = \sum_{r=0}^{\infty} a_r (r+k)(r+k-1) x^{r+k-2}$$

Tidy the O.D.E. and substitute in

$$2x^2 \frac{d^2y}{dx^2} + (2x^2 - x) \frac{dy}{dx} + y = 0$$

$$2x^2 \frac{d^2y}{dx^2} + 2x^2 \frac{dy}{dx} - x \frac{dy}{dx} + y = 0$$

$$\sum_{r=0}^{\infty} 2a_r (r+k)(r+k-1) x^{r+k} + \sum_{r=0}^{\infty} 2a_r (r+k) x^{r+k+1} - \sum_{r=0}^{\infty} a_r (r+k) x^{r+k} + \sum_{r=0}^{\infty} a_r x^{r+k} = 0$$

When $r=0$ the lowest power of x is x^k and the highest is x^{k+1}

Pull out the lowest power of x out of the summations

$$2a_0 k(k-1) x^k + \sum_{r=1}^{\infty} 2a_r (r+k)(r+k-1) x^{r+k} + \sum_{r=0}^{\infty} 2a_r (r+k) x^{r+k+1} - a_0 k x^k - \sum_{r=1}^{\infty} a_r (r+k) x^{r+k} + a_0 x^k + \sum_{r=1}^{\infty} a_r x^{r+k} = 0$$

Form an indicial equation from the expressions "pulled out" (lowest power)

$$\Rightarrow 2a_0 k(k-1) x^k - a_0 k x^k + a_0 x^k = 0$$

$$\Rightarrow [2k(k-1) - k + 1] a_0 x^k = 0$$

$$\Rightarrow [2k(k-1) - (k-1)] = 0 \quad a_0 \neq 0$$

$$\Rightarrow (k-1)(2k-1) = 0$$

$$\Rightarrow k = \begin{cases} 1 \\ \frac{1}{2} \end{cases}$$

DISTINCT ROOTS AND NOT DIFFERING BY AN INTEGER

NYB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 3

THE REST OF THE POWERS IN THE SUMMATIONS MUST ALSO EQUAL ZERO.

ADJUST THE SUMMATIONS SO THEY ALL START FROM $r=0$

$$\Rightarrow \sum_{r=1}^{\infty} 2a_r(r+k)(r+k-1)x^{r+k} + \sum_{r=0}^{\infty} 2a_r(r+k)x^{r+k+1} - \sum_{r=1}^{\infty} a_r(r+k)x^{r+k} + \sum_{r=1}^{\infty} a_r x^{r+k} = 0$$

$$\Rightarrow \sum_{r=0}^{\infty} 2a_{r+1}(r+k+1)(r+k)x^{r+k+1} + \sum_{r=0}^{\infty} 2a_r(r+k)x^{r+k+1} - \sum_{r=0}^{\infty} a_{r+1}(r+k+1)x^{r+k+1} + \sum_{r=0}^{\infty} a_{r+1}x^{r+k+1} = 0$$

$$\Rightarrow [2a_{r+1}(r+k+1)(r+k) + 2a_r(r+k) - a_{r+1}(r+k+1) + a_{r+1}]x^{r+k+1} = 0$$

$$\Rightarrow 2a_{r+1}(r+k+1)(r+k) - a_{r+1}(r+k+1) + a_{r+1} = -2a_r(r+k)$$

$$\Rightarrow [2(r+k+1)(r+k) - (r+k+1) + 1]a_{r+1} = -2a_r(r+k)$$

let $A = r+k$

$$\begin{aligned} 2(A+1)A - (A+1) + 1 &= 2A^2 + 2A - A - 1 + 1 \\ &= 2A^2 + A \\ &= A(2A+1) \end{aligned}$$

$$\Rightarrow (r+k)(2r+2k+1)a_{r+1} = -2a_r(r+k)$$

$$\Rightarrow (2r+2k+1)a_{r+1} = -2a_r$$

$$\Rightarrow a_{r+1} = -\frac{2}{2r+2k+1}a_r$$

NOW IF $k=1$ THIS RECURSIVE FORMULA BECOMES

$$a_{r+1} = -\frac{2}{2r+3}a_r$$

LYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 3

- IF $r=0$ $a_1 = -\frac{2}{3}a_0$
- IF $r=1$ $a_2 = -\frac{2}{5}a_1 = -\frac{2 \times 2}{3 \times 5}a_0$
- IF $r=2$ $a_3 = -\frac{2}{7}a_2 = -\frac{2 \times 2 \times 2}{3 \times 5 \times 7}a_0$
- IF $r=3$ $a_4 = -\frac{2}{9}a_3 = -\frac{2 \times 2 \times 2 \times 2}{3 \times 5 \times 7 \times 9}a_0$ ETC

THIS WE NOW HAVE IF $k=1$

$$\Rightarrow y_1 = a_0x + a_1x^2 + a_2x^3 + a_3x^4 + a_4x^5 + \dots$$

$$\Rightarrow y_1 = x[a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + \dots]$$

$$\Rightarrow y_1 = x\left[a_0 - \frac{2}{3}a_0x + \frac{2 \times 2}{3 \times 5}a_0x^2 - \frac{2 \times 2 \times 2}{3 \times 5 \times 7}a_0x^3 + \frac{2 \times 2 \times 2 \times 2}{3 \times 5 \times 7 \times 9}a_0x^4 + \dots\right]$$

$$\Rightarrow y_1 = a_0x\left[1 - \frac{2}{3}x + \frac{2^2}{3 \times 5}x^2 - \frac{2^3}{3 \times 5 \times 7}x^3 + \frac{2^4}{3 \times 5 \times 7 \times 9}x^4 + \dots\right]$$

$$\Rightarrow y_1 = a_0x\left[1 - \frac{2 \times 2}{3 \times 2}x + \frac{2^2 \times (4 \times 2)x^2}{5 \times 4 \times 3 \times 2} - \frac{2^3 \times (6 \times 4 \times 2)x^3}{7 \times 6 \times 5 \times 4 \times 3 \times 2} + \frac{2^4 \times (8 \times 6 \times 4 \times 2)x^4}{9 \times 8 \times 7 \times 6 \times 5 \times 4 \times 3 \times 2} + \dots\right]$$

$$\Rightarrow y_1 = a_0x\left[1 - \frac{2 \times 2 \times 1!}{3!}x + \frac{2^2 \times 2 \times 2!}{5!}x^2 - \frac{2^3 \times 2 \times 3!}{7!}x^3 + \frac{2^4 \times 2 \times 4!}{9!}x^4 + \dots\right]$$

$$\Rightarrow y_1 = a_0x\left[1 - \frac{2^2 \times 1!}{3!}x + \frac{2^4 \times 2!}{5!}x^2 - \frac{2^6 \times 3!}{7!}x^3 + \frac{2^8 \times 4!}{9!}x^4 + \dots\right]$$

$$\Rightarrow y_1 = a_0x \sum_{r=0}^{\infty} \left[\frac{2^{2r} \times r!}{(2r+1)!} (-1)^r x^r \right]$$

$$\Rightarrow y_1 = a_0x \sum_{r=0}^{\infty} \frac{4^r (-1)^r x^r r!}{(2r+1)!}$$

$$\Rightarrow y_1 = Ax \sum_{r=0}^{\infty} \frac{r!}{(2r+1)!} (-4x)^r$$

IVGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 3

NOW IF $l = \frac{1}{2}$ THE RECURSION EQUATION YIELDS

$$a_{r+1} = -\frac{a_r}{r+1}$$

- $r=0$ $a_1 = -a_0$
- $r=1$ $a_2 = -\frac{1}{2}a_1 = \frac{1}{2}a_0$
- $r=2$ $a_3 = -\frac{1}{3}a_2 = -\frac{1}{2 \times 3}a_0$
- $r=3$ $a_4 = -\frac{1}{4}a_3 = \frac{1}{2 \times 3 \times 4}a_0$ ETC

THUS WE NOW OBTAIN

$$y_2 = a_0 x^{\frac{1}{2}} + a_1 x^{\frac{3}{2}} + a_2 x^{\frac{5}{2}} + a_3 x^{\frac{7}{2}} + \dots$$

$$y_2 = x^{\frac{1}{2}} [a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots]$$

$$y_2 = x^{\frac{1}{2}} \left[a_0 - a_0 x + \frac{1}{2} a_0 x^2 - \frac{1}{2 \times 3} a_0 x^3 + \frac{1}{2 \times 3 \times 4} a_0 x^4 - \dots \right]$$

$$y_2 = a_0 x^{\frac{1}{2}} \left[1 - x + \frac{1}{2!} x^2 - \frac{1}{3!} x^3 + \frac{1}{4!} x^4 - \dots \right]$$

$$y_2 = B x^{\frac{1}{2}} e^{-x}$$

THUS THE GENERAL SOLUTION IS

$$y = y_1 + y_2 = A x \sum_{r=0}^{\infty} \frac{r!}{(r+1)!} (-4x)^r + B x^{\frac{1}{2}} e^{-x}$$

NYCB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 4

LOOKING AT $\mathcal{L}^{-1} \left[\ln \left(1 + \frac{1}{s^2} \right) \right]$ WE SUSPECT THAT THIS IS THE LAPLACE TRANSFORM OF $\frac{f(t)}{t}$ FOR SOME $f(t)$

$$\mathcal{L} \left[\frac{f(t)}{t} \right] = \int_s^\infty \bar{f}(x) dx = \bar{g}(s)$$

HERE $\bar{g}(s)$ IS $\ln \left(1 + \frac{1}{s^2} \right)$

$$\begin{aligned} \frac{d}{ds}(\bar{g}(s)) &= \frac{d}{ds} \left(\ln \left(1 + \frac{1}{s^2} \right) \right) = \frac{d}{ds} \left[\ln \left(\frac{s^2+1}{s^2} \right) \right] \\ &= \frac{d}{ds} \left[\ln(s^2+1) - \ln s^2 \right] = \frac{d}{ds} \left[\ln(s^2+1) - 2 \ln s \right] \\ &= \frac{2s'}{s^2+1} - \frac{2}{s} = 2 \left(\frac{s'}{s^2+1} \right) - 2 \left(\frac{1}{s} \right) \end{aligned}$$

WE RECOGNIZE THIS AS STANDARD RESULTS

$$\begin{aligned} 2 \mathcal{L} [\cos t] &= 2 \left(\frac{s'}{s^2+1} \right) \\ 2 \mathcal{L} [1] &= 2 \times \frac{1}{s} \end{aligned}$$

HENCE WE HAVE NOTING THE REVERSAL OF THE SIGNS DUE TO THE INTEGRATION LIMITS

$$\mathcal{L}^{-1} \left[\ln \left(1 + \frac{1}{s^2} \right) \right] = \frac{2(1 - \cos t)}{t}$$

QUICK CHECK

$$\begin{aligned} \mathcal{L} \left[\frac{2(1 - \cos t)}{t} \right] &= \int_s^\infty \mathcal{L} [2(1 - \cos t)] ds = \int_s^\infty \mathcal{L} [2 - 2 \cos t] ds = \int_s^\infty \left(\frac{2}{s} - \frac{2s'}{s^2+1} \right) ds \\ &= [2 \ln s - \ln(s^2+1)]_s^\infty = [\ln s^2 - \ln(s^2+1)]_s^\infty = \left[\ln \left(\frac{s^2}{s^2+1} \right) \right]_s^\infty \\ &= \cancel{\ln 1} - \ln \left(\frac{s^2}{s^2+1} \right) = -\ln \left(\frac{s^2}{s^2+1} \right) = \ln \left(\frac{s^2+1}{s^2} \right) = \ln \left(1 + \frac{1}{s^2} \right) \end{aligned}$$

YGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 5

START BY THE RECURSIVE PROPERTY OF THE GAMMA FUNCTION

$$\begin{aligned}
 \Gamma\left(n+\frac{1}{2}\right) &= \left(n-\frac{1}{2}\right)\left(n-\frac{3}{2}\right)\left(n-\frac{5}{2}\right)\dots \frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \frac{1}{2} \Gamma\left(\frac{1}{2}\right) \\
 &= \frac{1}{2}(2n-1) \times \frac{1}{2}(2n-3) \times \frac{1}{2}(2n-5) \dots \left(\frac{1}{2} \times 7\right)\left(\frac{1}{2} \times 5\right)\left(\frac{1}{2} \times 3\right)\left(\frac{1}{2} \times 1\right) \times \sqrt{\pi} \\
 &= \left(\frac{1}{2}\right)^n (2n-1)(2n-3)(2n-5) \dots 7 \times 5 \times 3 \times 1 \times \sqrt{\pi} \\
 &= \frac{1}{2^n} \frac{(2n-1)(2n-3)(2n-5) \dots 7 \times 5 \times 3 \times 1 \times \sqrt{\pi}}{(2n-2)(2n-4)(2n-6) \dots 6 \times 4 \times 2} \\
 &= \frac{1}{2^n} \times \frac{(2n-1)! \sqrt{\pi}}{2(n-1) \times 2(n-2) \times 2(n-3) \times \dots \times (2 \times 3) \times (2 \times 2) \times (2 \times 1)} \\
 &= \frac{1}{2^n} \times \frac{\Gamma(2n) \sqrt{\pi}}{2^{n-1} \times (n-1)(n-2)(n-3) \dots \times 3 \times 2 \times 1} \\
 &= \frac{1}{2^n} \times \frac{\Gamma(2n) \sqrt{\pi}}{2^{n-1} (n-1)!} \\
 &= \frac{\Gamma(2n) \sqrt{\pi}}{2^{2n-1} (n-1)!}
 \end{aligned}$$

As required

1YGB-MATHEMATICAL METHODS 3-PAPER C-QUESTION 6

STARTING WITH THE GENERATING FUNCTION

$$(1-2xt+t^2)^{-\frac{1}{2}} = \sum_{n=0}^{\infty} [t^n P_n(x)]$$

DIFFERENTIATE WITH RESPECT TO t

$$\Rightarrow -\frac{1}{2}(1-2xt+t^2)^{-\frac{3}{2}}(-2x+2t) = \sum_{n=0}^{\infty} [nt^{n-1} P_n(x)]$$

$$\Rightarrow (x-t)(1-2xt+t^2)^{-\frac{3}{2}} = \sum_{n=0}^{\infty} [nt^{n-1} P_n(x)]$$

$$\Rightarrow (x-t)(1-2xt+t^2)^{-\frac{3}{2}}(1-2xt+t^2) = (1-2xt+t^2) \sum_{n=0}^{\infty} [nt^{n-1} P_n(x)]$$

$$\Rightarrow (x-t)(1-2xt+t^2)^{-\frac{1}{2}} = (1-2xt+t^2) \sum_{n=0}^{\infty} [nt^{n-1} P_n(x)]$$

$$\Rightarrow (x-t) \sum_{n=0}^{\infty} [t^n P_n(x)] = (1-2xt+t^2) \sum_{n=0}^{\infty} [nt^{n-1} P_n(x)]$$

$$\Rightarrow \sum_{n=0}^{\infty} [t^n x P_n(x) - t^{n+1} P_n(x)] = \sum_{n=0}^{\infty} [nt^{n-1} P_n(x) - 2xnt^n P_n(x) + nt^{n+1} P_n(x)]$$

EQUATING COEFFICIENTS OF t, SAY [t^n]

$$\Rightarrow x P_n(x) - P_{n+1}(x) = (n+1) P_{n+1}(x) - 2x n P_n(x) + (n-1) P_{n-1}(x)$$

$$\Rightarrow 0 = (n+1) P_{n+1}(x) - 2x n P_n(x) - x P_n(x) + (n-1) P_{n-1}(x) + P_{n+1}(x)$$

$$\Rightarrow 0 = (n+1) P_{n+1}(x) - (2x n + x) P_n(x) + n P_{n-1}(x)$$

$$\Rightarrow (n+1) P_{n+1}(x) - (2n+1)x P_n(x) + n P_{n-1}(x) = 0$$

AS REQUIRED

NOB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 7

START WITH A SUBSTITUTION FIRST

$$\begin{aligned} u = \frac{1}{x+1} &\Leftrightarrow x+1 = \frac{1}{u} \quad \left(x = \frac{1}{u} - 1\right) \\ &\Rightarrow dx = -\frac{1}{u^2} du \\ x=0 &\mapsto u=1 \\ x=\infty &\mapsto u=0 \end{aligned}$$

TRANSFORMING THE INTEGRAL INCLUDING LIMITS

$$\begin{aligned} \int_1^0 \frac{1}{\sqrt{\frac{1}{u}-1}} \cdot \frac{1}{u} \left(-\frac{1}{u^2} du\right) &= \int_1^0 \frac{-1}{\sqrt{\frac{1-u}{u}} \cdot \frac{1}{u} u^2} du \\ &= \int_0^1 \frac{+1}{\frac{\sqrt{1-u}}{\sqrt{u}} \times u} du \\ &= \int_0^1 \frac{1}{(1-u)^{\frac{1}{2}} (u^{\frac{1}{2}})} du \\ &= \int_0^1 (1-u)^{-\frac{1}{2}} u^{-\frac{1}{2}} du \end{aligned}$$

NOW USING THE BETA FUNCTION DEFINITION

$$\dots = \int_0^1 (1-u)^{\frac{1}{2}-1} u^{\frac{1}{2}-1} du = B\left(\frac{1}{2}, \frac{1}{2}\right)$$

SWITCHING INTO GAMMA FUNCTIONS FOR THE FINAL ANSWER

$$= \frac{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2}\right)}{\Gamma(1)} = \frac{\sqrt{\pi} \sqrt{\pi}}{0!} = \frac{\pi}{1} = \underline{\underline{\pi}}$$

1YGB - MATHEMATICAL METHODS 3 - PART C - QUESTION 8

IT IS BEST TO WORK WITH EXPONENTIALS IN THIS QUESTION

$$f(z) = \frac{ze^{kz}}{z^4+1} \text{ HAS SIMPLE POLES AT:}$$

$$z^4 = -1 = e^{i(\pi+2n\pi)} = e^{i\pi(2n+1)}, \quad n=0,1,2,3$$

$$z = e^{i\frac{\pi}{4}(2n+1)}, \quad n=0,1,2,3 \quad (\text{or } -2, -1, 0, 1)$$

$$z = e^{i\frac{\pi}{4}}, e^{i\frac{3\pi}{4}}, e^{-i\frac{\pi}{4}}, e^{-i\frac{3\pi}{4}}$$

CALCULATE THESE RESIDUES USING A GENERAL METHOD - LET A POLE BE AT $z=z_0$

$$\text{Res}(f; z_0) = \lim_{z \rightarrow z_0} \left[\frac{(z-z_0)ze^{kz}}{z^4+1} \right]$$

THIS MUST PRODUCE 'ZERO OVER ZERO' AS $z-z_0$ MUST BE A FACTOR OF THE DENOMINATOR SO WE PROCEED BY L'HOSPITAL RULE

$$\text{Res}(f; z_0) = \lim_{z \rightarrow z_0} \left[\frac{\frac{d}{dz}[(z-z_0)ze^{kz}]}{\frac{d}{dz}(z^4+1)} \right] = \lim_{z \rightarrow z_0} \left[\frac{ze^{kz} + (z-z_0)e^{kz} + (z-z_0)ke^{kz}}{4z^3} \right]$$

$$\text{Res}(f; z_0) = \frac{z_0 e^{kz_0}}{4z_0^3} = \boxed{\frac{e^{kz_0}}{4z_0^2}}$$

NEXT OBTAIN THE RESIDUE AT EACH OF THE FOUR POLES

$$z_0 = e^{i\frac{\pi}{4}} \text{ GIVES } \frac{e^{ke^{i\frac{\pi}{4}}}}{4e^{i\frac{\pi}{2}}} = \frac{e^{k\left(\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2}\right)}}{4i}$$

$$z_0 = e^{-i\frac{\pi}{4}} \text{ GIVES } \frac{e^{ke^{-i\frac{\pi}{4}}}}{4e^{-i\frac{\pi}{2}}} = \frac{e^{k\left(\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}\right)}}{-4i}$$

$$z_0 = e^{i\frac{3\pi}{4}} \text{ GIVES } \frac{e^{ke^{i\frac{3\pi}{4}}}}{4e^{i\frac{3\pi}{2}}} = \frac{e^{k\left(-\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2}\right)}}{-4i}$$

$$z_0 = e^{-i\frac{3\pi}{4}} \text{ GIVES } \frac{e^{ke^{-i\frac{3\pi}{4}}}}{4e^{-i\frac{3\pi}{2}}} = \frac{e^{k\left(-\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2}\right)}}{4i}$$

IYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 8

ADDING THE 4 RESIDUES - LET $a = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$ FOR SIMPLICITY

$$\text{SUM OF 4 RESIDUES} = \frac{e^{k(\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2})}}{4i} + \frac{e^{k(\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2})}}{-4i} + \frac{e^{k(-\frac{\sqrt{2}}{2} + i\frac{\sqrt{2}}{2})}}{-4i} + \frac{e^{k(-\frac{\sqrt{2}}{2} - i\frac{\sqrt{2}}{2})}}{4i}$$

$$= \frac{1}{4i} \left[e^{k(a+ai)} - e^{k(a-ai)} - e^{k(-a+ai)} + e^{k(-a-ai)} \right]$$

$$= \frac{1}{4i} \left[e^{ak} e^{kai} - e^{ak} e^{-kai} + e^{-ak} e^{-iak} - e^{-ak} e^{aki} \right]$$

$$= \frac{1}{4i} \left[e^{ak} (e^{aki} - e^{-aki}) - e^{-ak} (e^{aki} - e^{-aki}) \right]$$

$$= \frac{1}{4i} \left[(e^{aki} - e^{-aki}) (e^{ak} - e^{-ak}) \right]$$

$$= \frac{1}{4i} \times 2\sinh(aki) \times 2\sinh(ak)$$

$$= \cancel{\frac{1}{4i}} \times \cancel{2i} \sin(ak) \times \cancel{2}\sinh(ak)$$

$$= \sin(ak) \sinh(ak)$$

$$= \sin\left(\frac{k}{\sqrt{2}}\right) \sinh\left(\frac{k}{\sqrt{2}}\right)$$

As required

LYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 9

STARTING WITH THE GENERATING FUNCTION FOR THE BESSEL FUNCTIONS OF THE FIRST KIND

$$\Rightarrow e^{\frac{1}{2}x(t+\frac{1}{t})} = \sum_{n=-\infty}^{\infty} [t^n J_n(x)]$$

$$\Rightarrow e^{\frac{1}{2}x(t+\frac{1}{t})} e^{\frac{1}{2}y(t+\frac{1}{t})} = \left[\sum_{k=-\infty}^{\infty} [t^k J_k(x)] \right] \left[\sum_{m=-\infty}^{\infty} [t^m J_m(y)] \right]$$

$$\Rightarrow e^{\frac{1}{2}(x+y)(t+\frac{1}{t})} = \sum_{k=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} [J_k(x) J_m(y) t^{k+m}]$$

LET $n = k+m$ AND NOTE THAT SUMMATION UNITS ARE UNCHANGED

$$\Rightarrow e^{\frac{1}{2}(x+y)(t+\frac{1}{t})} = \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} [t^n J_{n-m}(x) J_m(y)]$$

$$\Rightarrow e^{\frac{1}{2}(x+y)(t+\frac{1}{t})} = \sum_{n=-\infty}^{\infty} [t^n \sum_{m=-\infty}^{\infty} [J_m(y) J_{n-m}(x)]]$$

$$\Rightarrow \sum_{n=-\infty}^{\infty} [t^n J_n(x+y)] = \sum_{n=-\infty}^{\infty} [t^n \left[\sum_{m=-\infty}^{\infty} [J_m(y) J_{n-m}(x)] \right]]$$

$$\Rightarrow \underline{J_n(x+y) = \sum_{m=-\infty}^{\infty} [J_m(y) J_{n-m}(x)]}$$


AS REQUIRED

- 1 -

LYGB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 10

- START WITH THE DIAGRAM OPPOSITE

$$x = 8\cos^3 t, \quad y = \sin^3 t, \quad 0 \leq t \leq \frac{\pi}{2}$$



- THE MASS OF THE INFINITESIMAL DISC OF RADIUS y & THICKNESS δx IS GIVEN BY

$$\delta m = \pi \rho y^2 \delta x \quad (\rho = \text{DENSITY})$$

- THE "MOMENT" OF THIS INFINITESIMAL MASS, ABOUT THE y AXIS IS GIVEN BY

$$x \delta m = x (\pi \rho y^2 \delta x) = \pi \rho y^2 x \delta x$$

- SUMMING UP, TAKING LIMITS, WE OBTAIN

$$\Rightarrow M \bar{x} = \int_{x=0}^8 \pi \rho y^2 x \, dx$$

$$\Rightarrow \bar{x} \int_{x=0}^8 \pi \rho y^2 \, dx = \int_{x=0}^8 \pi \rho y^2 x \, dx$$

$$\Rightarrow \bar{x} \int_{t=\pi/2}^0 (\sin^3 t)^2 \underbrace{(-24 \cos^2 t \sin t \, dt)}_{dx} = \int_{t=\pi/2}^0 (\sin^3 t)^2 \underbrace{(-24 \cos^2 t \sin t \, dt)}_{dx} \underbrace{(8 \cos^3 t)}_x$$

$$\Rightarrow \bar{x} \int_0^{\pi/2} \sin^7 t \cos^2 t \, dt = \int_0^{\pi/2} 8 \sin^7 t \cos^5 t \, dt$$

LYGB - MATHEMATICAL METHODS 3 - QUESTION 10

● EVALUATE USING BETA FUNCTIONS

$$\Rightarrow \bar{x} \int_0^{\frac{\pi}{2}} 2 (\sin t)^{2 \times 4 - 1} (\cos t)^{2 \times \frac{3}{2} - 1} dt = 8 \int_0^{\frac{\pi}{2}} 2 (\sin t)^{2 \times 4 - 1} (\cos t)^{2 \times 3 - 1} dt$$

$$\Rightarrow \bar{x} B(4, \frac{3}{2}) = 8 B(4, 3)$$

$$\Rightarrow \bar{x} \frac{\Gamma(4) \Gamma(\frac{3}{2})}{\Gamma(\frac{11}{2})} = \frac{8 \Gamma(4) \Gamma(3)}{\Gamma(7)}$$

$$\Rightarrow \bar{x} = \frac{8 \Gamma(3) \Gamma(\frac{11}{2})}{\Gamma(7) \Gamma(\frac{3}{2})}$$

$$\Rightarrow \bar{x} = \frac{8 \times 2!}{6!} \times \frac{\frac{9}{2} \times \frac{7}{2} \times \frac{5}{2} \times \frac{3}{2} \times \Gamma(\frac{3}{2})}{\Gamma(\frac{3}{2})}$$


$$\Rightarrow \bar{x} = \frac{\cancel{16}}{720} \times \frac{9 \times 7 \times 5 \times 3}{\cancel{16}}$$

$$\Rightarrow \bar{x} = \frac{21}{16}$$

YGB - MATHEMATICAL METHODS 3 - PAGE 2 C-QUESTION 11

BY THE INTEGRAL DEFINITION OF FOURIER TRANSFORM

$$f(x) = \frac{1}{(a^2 + x^2)^2}, \quad a > 0$$




$$\hat{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ikx} dx$$

$$\hat{f}(k) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} \frac{\cos kx - i \sin kx}{(a^2 + x^2)^2} dx$$

$$\hat{f}(k) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{\cos kx}{(a^2 + x^2)^2} dx$$

ONLY EVEN PART SURVIVES

PROCEED BY CONTOUR INTEGRATION - USE DIFFERENT CONTOUR DEPENDING ON THE SIGN OF k



$$f(z) = \frac{e^{ikz}}{(z^2 + a^2)^2} \text{ HAS DOUBLE}$$

POLES AT $\pm ai$, ONE IN EACH OF THE CONTOURS

SHOWN OPPOSITE

RESIDUE AT $z = ai$

$$\begin{aligned} \lim_{z \rightarrow ai} \left[\frac{d}{dz} \left[f(z) (z - ai)^2 \right] \right] &= \lim_{z \rightarrow ai} \left[\frac{d}{dz} \left[\frac{e^{ikz} \cancel{(z - ai)^2}}{\cancel{(z - ai)^2} (z + ai)^2} \right] \right] \\ &= \lim_{z \rightarrow ai} \left[\frac{d}{dz} \left(\frac{e^{ikz}}{(z + ai)^2} \right) \right] \end{aligned}$$

1YGB-MATHEMATICAL METHODS 3 - PAPER C - QUESTION 11

$$\begin{aligned}
 &= \lim_{z \rightarrow ai} \left[\frac{ike^{ikz}(z+ai)^2 - 2(z+ai)e^{ikz}}{(z+ai)^4} \right] \\
 &= \lim_{z \rightarrow ai} \left[\frac{ike^{ikz}(z+ai) - 2e^{ikz}}{(z+ai)^3} \right] = \lim_{z \rightarrow ai} \left[\frac{e^{ikz}(ikz - ak - 2)}{(z+ai)^3} \right] \\
 &= \frac{e^{-ak}(-2ak - 2)}{(2ai)^3} = \frac{-2e^{-ak}(ak+1)}{-8a^3i} = \frac{(ak+1)e^{-ak}}{4a^3i}
 \end{aligned}$$

IN A MOST ANALOGOUS FASHION THE RESIDUE AT $-ai$ FOR USE IN THE BOTTOM CONTOUR

$$\begin{aligned}
 \dots &= \lim_{z \rightarrow -ai} \left[\frac{ike^{ikz}(z-ai) - 2e^{ikz}}{(z-ai)^3} \right] \\
 &= \lim_{z \rightarrow -ai} \left[\frac{e^{ikz}(ikz + ak - 2)}{(z-ai)^3} \right] = \frac{e^{ak}(2ak - 2)}{(-2ai)^3} \\
 &= \frac{2(ak-1)e^{ak}}{8a^3i} = \frac{(ak-1)e^{ak}}{4a^3i}
 \end{aligned}$$

• IF $k > 0$ USING THE "TOP" CONTOUR

$$\int_{\Gamma_1} f(z) dz = \int_{-R}^R \frac{e^{ikx}}{(x^2+a^2)^2} dx + \int_{\gamma_1} \frac{e^{ikz}}{(z^2+a^2)^2} dz = 2\pi i \left(\frac{(ak+1)e^{-ak}}{4a^3i} \right)$$

• IF $k < 0$ USING THE "BOTTOM" CONTOUR

$$\int_{\Gamma_2} f(z) dz = \int_{-R}^R \frac{e^{ikx}}{(x^2+a^2)^2} dx + \int_{\gamma_2} \frac{e^{ikz}}{(z^2+a^2)^2} dz = \underset{\substack{\uparrow \\ \text{CLOCKWISE PATH}}}{-2\pi i} \left(\frac{(ak-1)e^{ak}}{4a^3i} \right)$$

THE INTEGRALS OVER THE ARCS γ_1 & γ_2 VANISH AS $R \rightarrow \infty$, AS BOTH SATISFY THE CONDITIONS OF JORDAN'S LEMMA, FOR THE CORRECT SIGN OF k IN EACH CONTOUR

NYSB - MATHEMATICAL METHODS 3 - PAPER C - QUESTION 11

DEALING WITH EACH CASE SEPARATELY AS $R \rightarrow \infty$

• IF $k > 0$

$$\int_{-\infty}^{\infty} \frac{e^{ikx}}{(x^2+a^2)^2} dx = \frac{\pi}{2a^3} (ak+1) e^{-ak}$$

$$\int_0^{\infty} \frac{2\cos kx}{(x^2+a^2)^2} dx = \frac{\pi e^{-ak} (ak+1)}{2a^3}$$

$$\sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{2\cos kx}{(x^2+a^2)^2} dx = \sqrt{\frac{\pi}{2}} \frac{e^{-ak} (ak+1)}{a^3}$$

$$\sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{\cos kx}{(x^2+a^2)^2} dx = \sqrt{\frac{\pi}{8}} \frac{e^{-ak} (ak+1)}{a^3}$$

• IF $k < 0$

$$\int_{-\infty}^{\infty} \frac{e^{ikx}}{(x^2+a^2)^2} dx = \frac{\pi}{2a^3} (-ak+1) e^{ak}$$

$$\sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{\cos kx}{(x^2+a^2)^2} dx = \sqrt{\frac{\pi}{8}} \frac{e^{ak} (-ak+1)}{a^3}$$

COMBINING THE RESULTS FOR ALL k

$$\hat{f}(k) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{\cos kx}{(x^2+a^2)^2} dx = \begin{cases} \sqrt{\frac{\pi}{8}} \frac{e^{-ak} (ak+1)}{a^3} & k > 0 \\ \sqrt{\frac{\pi}{8}} \frac{e^{ak} (-ak+1)}{a^3} & k < 0 \end{cases}$$

$$\hat{f}(k) = \sqrt{\frac{\pi}{8}} \frac{(1+|k|a) e^{-|k|a}}{a^3}$$